

AD-A059 022

TEMPLE UNIV PHILADELPHIA PA DEPT OF MATHEMATICS
STATISTICAL ASPECTS OF EXTREME VALUE THEORY.(U)
JUL 78 J GALAMBOS

F/6 12/1

UNCLASSIFIED

AFOSR-TR-78-1139

AFOSR-78-3504

ML

| OF |
AD
4059022



END
DATE
FILMED
11-78
DDC

AD A0 59022
DDC FILE COPY

LEVEL II

2

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

19 REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR-TR-78-1139	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) STATISTICAL ASPECTS OF EXTREME VALUE THEORY.	5. TYPE OF REPORT & PERIOD COVERED Interim rept.	
7. AUTHOR(s) Janos Galambos	8. CONTRACT OR GRANT NUMBER(s) AFOSR-78-3504	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Temple University Department of Mathematics Philadelphia, PA 19122	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2304/A5	
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research/NM Boiling AFB, Washington, DC 20332	12. REPORT DATE July 1978	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. NUMBER OF PAGES 5	
	15. SECURITY CLASS. (of this report) UNCLASSIFIED	
15a. DECLASSIFICATION DOWNGRADING SCHEDULE		
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of abstract entered in Block 20, if different from Report) D		
18. SUPPLEMENTARY NOTES Presented at the Australian Statistical Conference, 11-14 July 1978, Canberra, Australia		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Let X_1, X_2, \dots, X_n be random variables with common distribution function $F(x)$. Put $Z_n = \max(X_1, X_2, \dots, X_n)$. Assume that the interdependence of the X_j is such that, with suitable constants a_n and $b_n > 0$, $P(Z_n < a_n + b_n x)$ is asymptotically $F^n(a_n + b_n x)$; see Chapter 3 in the book by the present author,		

20. Abstract

J. Galambos (1978) for models with this property. Hence, for most functions $F(x)$ occurring in practice, $(Z_n - a_n)/b_n$ has an asymptotic distribution $H(x)$ as $n \rightarrow +\infty$. This asymptotic distribution is necessarily one of the three classical types obtained in the 1920s by Fisher, Tippet and Frechet. However, if $F(x)$ is not known (which is the case of most practical situations), then neither the normalizing constants a_n and b_n , nor the actual type of the limit distribution $H(x)$ can be determined. Since goodness of fit tests are not sensitive for the tail of the distribution $F(x)$, and since an estimate for $F(x)$ is not suitable in extreme value theory (see Galambos (1978), Example 2.6.3), some other techniques are needed which are free from $F(x)$. The paper is devoted to such a technique, which is an extension of the approach by J. Pickands III (1975). In such a method, one uses a number of upper extremes and the parametric form of the three classical types for $H(x)$, known as the von Mises form.

ACCESSION NO.	
DTIC	White Section <input checked="" type="checkbox"/>
DDI	Defi Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION.....	
BY.....	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SPECIAL
A	

Statistical aspects of extreme value theory


by

Janos Galambos

Department of Mathematics
Temple University
Philadelphia, Pa. 19122.

Preliminary Report. Presented at the
Australian Statistical Conference,
11-14 July, 1978, Canberra, Australia.

Research supported by a Grant (#78-3504) from AFOSR
to Temple University.



78 07 20 094

A b s t r a c t

Let X_1, X_2, \dots, X_n be random variables with common distribution function $F(x)$. Put $Z_n = \max(X_1, X_2, \dots, X_n)$. Assume that the interdependence of the X_j is such that, with suitable constants a_n and $b_n > 0$, $P(Z_n < a_n + b_n x)$ is asymptotically $F^n(a_n + b_n x)$; see Chapter 3 in the book by the present author, J. Galambos (1978) for models with this property. Hence, for most functions $F(x)$ occurring in practice, $(Z_n - a_n)/b_n$ has an asymptotic distribution $H(x)$ as $n \rightarrow +\infty$. This asymptotic distribution is necessarily one of the three classical types obtained in the 1920s by Fisher, Tippet and Frechet. However, if $F(x)$ is not known (which is the case of most practical situations), then neither the normalizing constants a_n and b_n , nor the actual type of the limit distribution $H(x)$ can be determined. Since goodness of fit tests are not sensitive for the tail of the distribution $F(x)$, and since an estimate for $F(x)$ is not suitable in extreme value theory (see Galambos (1978), Example 2.6.3), some other techniques are needed which are free from $F(x)$. The paper is devoted to such a technique, which is an extension of the approach by J. Pickands III (1975). In such a method, one uses a number of upper extremes and the parametric form of the three classical types for $H(x)$, known as the von Mises form.

References

- Galambos, Janos (1978). The asymptotic theory of extreme order statistics. John Wiley and Sons, New York.
- Pickands, James III (1975). Statistical inference using extreme order statistics. *Ann. Statist.* 3, 119-131.

78 07 20 094

A new statistical method

Let X_1, X_2, \dots, X_n be identically distributed random variables. Put $Z_n = \max(X_1, X_2, \dots, X_n)$. Assume that there are constants a_n and $b_n > 0$ such that $P(Z_n < a_n + b_n x)$ can be approximated by $F^n(a_n + b_n x)$, where F is usually the common distribution function of the X 's. This implies that if F has a smooth enough tail, then $(Z_n - a_n)/b_n$ has an asymptotic distribution $H(x)$. Our aim is to determine the form of $H(x)$ in the case when F is not known. The approximation of $F(x)$ by the empirical distribution function, or by other classical methods, is inefficient in extreme value theory; see Galambos (1978a) for detailed discussion. Hence a direct method is required to get information on $H(x)$ without trying to determine F . There is only one work on this line in the literature by J. Pickands III (1975). Here we consider an alternate model. While Pickands develops a method, in which data will automatically separate the form of $H(x)$ out of the three possibilities, our method is a hypothesis test, in which we make a decision on the mentioned form. It remains an unsolved problem to compare the two methods on a theoretical basis. Because the nature of the two approaches is different, a clear comparison is difficult. However, advantages and disadvantages of one method or the other can be clarified from several points of view. As a matter of opinion, we can state that the method by Pickands would be preferable to a test if the accuracy of his method were available. Since the Pickands method involves three different passages to limit for supporting its conclusions, an estimate of accuracy is quite important. On the other hand, in some cases, one would need the actual form of $H(x)$, for which a test is suitable, while the method of Pickands leaves this undecided.

The method described below is in an early stage of investigation. However, initial results and some numerical computations suggest that it is a useful attempt and it thus deserves a thorough investigation.

Basic assumptions: 1./ The X_j are identically distributed with common distribution function $F(x)$; 2./ The dependence of the X_j is such that the distribution of Z_n can be approximated by the n -th power of $F(x)$ for large values of x ; 3./ $F(x) < 1$ for all x ; 4./ $E(X_j)$ is finite; and 5./ There are constants a_n and $b_n > 0$ such that $(Z_n - a_n)/b_n$ has an asymptotic distribution $H(x)$.

The above assumptions imply that there are constants A and $B > 0$ such that $H(A+Bx)$ equals either $H_{3,0}(x) = \exp(-e^{-x})$ or $H_{1,c}(x) = \exp(-x^{-c})$, $x > 0$, $c > 0$. Our aim is to decide by a statistical method based on the X_j that which of the two forms mentioned applies.

A test for the form of $H(x)$. Notice that the X_j provide a single observation on Z_n . Yet a number of extremes can be used to provide information on $H(x)$. The actual method is as follows. Choose a number t . Select those X_j that exceed t . Transform such X_j 's to

$$Y_j = (X_j - t) / E(X_j - t \mid X_j > t) .$$

Test that the distribution of these Y_j is unit exponential against the composite hypothesis that this distribution belongs to the Pareto family. Accepting the hypothesis means acceptance of $H(A+Bx) = H_{3,0}(x)$, while rejection of the null hypothesis leads to the acceptance of $H_{1,c}(x)$ as the limiting distribution of the properly normalized maximum. The power of the test depends on the choice of t , which choice is yet to be investigated in terms of some optimality criterion.

The theoretical justification of the above procedure is a classical criterion, combined with a result of L. de Haan (1970), for distributions belonging to given domains of attraction for the maxima. Details will be given in an expanded form of this present paper.

3. References

- Galambos, J. (1978). The asymptotic theory of extreme order statistics. John Wiley and Sons, New York.
- Galambos, J. (1978a). Extreme value theory in applied probability. Preliminary report.
- Haan, L. de (1970). On regular variation and its application to the weak convergence of sample extremes. Mathematical Centre Tracts, Vol. 32, Amsterdam.
- Pickands, J. III (1975). Statistical inference using extreme order statistics. Ann. Statist. 3, 119-131.